

**Note - ibuprofen and acetaminophen may be dosed alternately in infants over 6 mos of age every 4 hours*

**ACETAMINOPHEN
(TYLENOL)**

dosed every 4 hrs

Infant's
Concentrated Drops
80 mg/0.8 mL
Dropperful
(not widely available)

Infant's/Children's
Suspension Liquid
160 mg/5 ml

Children's Soft
Chewable Tablets
80 mg Tablet

Junior Strength
Chewable Tablets
160mg Tablet

Weight

Dropper

**Dosing Cup or
syringe**

Tablet

Tablet

6-11 lbs

1/2 (0.4mL)

ask MD

12-17 lbs

1=(0.8mL)

1/2 tsp.

18-23 lbs

1-1/2=(1.2 mL)

3/4 tsp.

24-35 lbs

2=(1.6mL)

1 tsp.

2

36-47 lbs

1-1/2 tsp.

3

48-59 lbs

2 tsp.

4

2

60-71 lbs

2-1/2 tsp.

5

2-1/2

72-95 lbs

3 tsp.

6

3

96 lbs & over

4 tsp.

8

4

**Note - ibuprofen and acetaminophen may be dosed alternately in infants over 6 mos of age every 4 hours*

IBUPROFEN
(Advil, Motrin)

not recommended for infants under 6 mos

dosed every 6 hrs

Infant's
Ibuprofen
Concentrated
Drops
50mg/1.25mL

Children's
Ibuprofen
Suspension
Tablets
100mg/5mL

Children's
Soft
Chewable
Tablets
50mg each

Junior
Strength
Chewable
Caplets
100mg each

Junior
Strength
Ibuprofen
100mg

Weight

Dropper

Teaspoon

Tablet

Tablet

Caplet

6-11 lbs

**Not
Recommended**

12-17 lbs

1=(1.25mL)

1/2 tsp.

18-23 lbs

1-1/2=(1.875mL)

3/4 tsp.

24-35 lbs

2=(2.5mL)

1 tsp.

2

36-47 lbs

1-1/2 tsp.

3

48-59 lbs

2 tsp.

4

2

2

60-71 lbs

2-1/2 tsp.

5

2-1/2

2-1/2

72-95 lbs

3 tsp.

6

3

3

96 lbs & over

4 tsp.

8

4

4